



Domain reporting in Systemic Sclerosis-Related Digital Ulcers: An OMERACT Scoping Review

Michael Hughes^{a,b,*}, Nancy Maltez^c, Edith Brown^d, Virginia Hickey^e, Beverley Shea^c, John D Pauling^{f,g}, Susanna Proudman^h, Peter A. Merkelⁱ, Ariane L. Herrick^b

^a Northern Care Alliance NHS Foundation Trust, Salford Care Organisation, Salford, UK

^b Division of Musculoskeletal and Dermatological Sciences, The University of Manchester, Manchester Academic Health Science Centre, Manchester, UK

^c Faculty of Medicine, University of Ottawa, Ottawa, Ontario, Canada

^d Manchester, United Kingdom

^e Adelaide, Australia

^f North Bristol NHS Trust, Bristol, UK

^g Bristol Medical School, University of Bristol, Bristol, UK

^h Discipline of Medicine, University of Adelaide and Rheumatology Unit, Royal Adelaide Hospital, Adelaide, Australia

ⁱ Division of Rheumatology, Department of Medicine, Division of Epidemiology, Department of Biostatistics, Epidemiology, and Informatics, University of Pennsylvania, Philadelphia, PA, USA

ARTICLE INFO

Keywords:

Systemic sclerosis
Scleroderma
Digital ulcers
Outcomes
Clinical Trials
Domains, OMERACT, Scoping review

ABSTRACT

Background: Digital ulcers (DUs) are a major cause of pain and disability in patients with systemic sclerosis (SSc). The aim of this scoping review was to evaluate the outcome domains used in studies of SSc-associated DUs.

Methods: Electronic databases (EMBASE, MEDLINE and the Cochrane Library) were searched for articles written (1947 onwards) in English relating to SSc-DUs. A minimum of 15 participants for studies of imaging and 25 participants for questionnaire-based studies was required for inclusion. Information on all primary and secondary domains was extracted.

Results: 4869 manuscripts were identified, of which 40 met the eligibility criteria and were included in the synthesis. Most studies were randomized controlled trials (n=13), or prospective (n=12)/retrospective (n=8) observational studies. Interventions included oral or intravenous drugs (n=25), topical/local treatments (n=5), and surgical interventions (n=2). Approximately half the studies assessed either the count/number of DUs (n=23) and/or improvement in DUs (n=20). Functional impact of DUs was examined in 25% (n=10) of studies. Other domains were related to complications of DUs (n=7), pain (n=6), health-related quality of life (n=4), microvascular assessment/pathophysiology (n=4), global assessment of DUs (n=2), and histopathology (n=1).

Conclusion: This scoping review identified a broad range of disease-related domains used to study SSc-DUs. There is significant heterogeneity in these domains. These data will inform the ongoing work of the OMERACT Vascular Disease in Systemic Sclerosis Working Group to define a core set of disease broad domains to capture the burden of DUs in SSc.

Introduction

Digital ulcers (DUs) occur in over half of patients with systemic sclerosis (SSc) and are associated with significant pain and disability

[1–3]. DUs are often defined as lesions with a loss of surface epithelization and discernible depth with a break in the basement membrane [4, 5]. DUs in SSc often occur early (within the first 5 years) in the course of the disease [1,6,7]. The pathogenesis of DUs is incompletely understood,

Abbreviations: DUs, Digital ulcers; SSc, Systemic sclerosis.

* Corresponding author: Dr Michael Hughes, Consultant Rheumatologist. Department of Rheumatology. Northern Care Alliance NHS Foundation Trust, Salford Care Organisation, Salford, UK

** Honorary Senior Clinical Lecturer. Division of Musculoskeletal and Dermatological Sciences. The University of Manchester, Manchester, UK, Telephone: 0161 922 6000

E-mail address: Michael.hughes-6@postgrad.manchester.ac.uk (M. Hughes).

<https://doi.org/10.1016/j.semarthrit.2023.152220>

Available online 5 May 2023

0049-0172/© 2023 Elsevier Inc. All rights reserved.

although ischemia is believed to play a central role, particularly in fingertip ulcers [3,6,8]. Mechanical factors also likely drive ulceration on the dorsal aspects of the hands (e.g., overlying the small joints), an area vulnerable to recurrent microtrauma [8,9]. DUs can also develop in relation to underlying subcutaneous calcinosis [10,11]. Complications of ulcers, including infection and gangrene, can significantly delay healing and may require surgical intervention [2,12–14].

DUs are also associated with a severe disease course of SSc, including internal organ involvement [15,16]. Despite current treatment approaches, including drug therapies, DUs are often slow to heal, and many patients experience recurrent digital ulceration [9,17]. Pain is a cardinal feature of DUs, and patients also often experience other intrusive physical symptoms (e.g., sensitivity) [18–20]. DUs are associated with significant emotional impact, including effects on personal relationships, and impairment of physical, social, and occupational activities [19]. Furthermore, there is a significant societal burden from SSc-DUs, including the costs of healthcare utilization [21].

Assessment of DU burden (impact and severity) is challenging in both clinical practice and clinical trials, posing a major barrier to the development of new and optimized treatment approaches for DUs, including non-pharmacological interventions. Agreement among experts in SSc assessing DUs is poor to moderate [5,22–24]. Furthermore, agreement between patients' and rheumatologists' assessments is also poor, even when clinicians are aware of 'real-world' clinical contextual information, such as the presence of discharge [24]. Much of the current understanding of the impact of SSc-DUs is derived from cross-sectional studies utilizing patient-reported outcome instruments to assess broader aspects of SSc disease severity [18].

Against this background, this scoping review of the literature aimed to evaluate the broad domains of illness, and the range of instruments and outcome domains, used in clinical studies of DUs in patients with SSc.

Methods

Working group

This scoping review [25], was conducted by the Outcome Measures in Rheumatology (OMERACT) Vascular Disease in Systemic Sclerosis Working Group which consists of six clinicians with an interest in SSc-DUs, a methodologist, and two patient research partners [26]. This project followed the OMERACT domain selection process [27].

Search strategy

A literature search strategy (**Supplementary Material**) was developed for use in EMBASE (OVID interface, 1947 onwards), MEDLINE (OVID interface, 1947 onwards), and Cochrane Central Register of Controlled Trials (OVID interface, 1947 onwards). These databases were searched for studies pertaining to participants with SSc-DUs. The databases were searched for studies pertaining to participants with a clinical diagnosis of SSc-associated DU with no limitation by classification criteria used, (given the various iterations in classification criteria for SSc utilized over the study period). Furthermore, patients were not limited to having a DU at baseline, but were required to have undergone an assessment for DUs.

Eligibility criteria

There was no limitation by intervention, comparator, or study setting. Randomized control trials (RCTs), quasi-randomized studies, case-control studies, prospective and retrospective cohort studies, case series, and cross-sectional studies written in English were eligible. Due to the large number of studies examining SSc-DU, this analysis was limited to studies with a minimum of 15 participants for studies of imaging modalities and 25 participants for questionnaire-based studies

were required for inclusion. Basic laboratory, genetic, or pre-clinical studies, and articles that were only available in abstract form were excluded.

Data extraction

Scoping review sources were uploaded to a citation management software (Covidence) and duplicate files were deleted. Two authors (MH, NM) independently completed screening of the title/abstract body and full text according to the inclusion criteria outlined above. Disagreements were resolved through consensus between the screening authors. A standardized data extraction form was developed and approved by all the study authors. Data extraction was independently piloted by two review authors (MH, NM) by extracting pertinent data for the first ten studies deemed eligible for inclusion. Thereafter, the remainder of the data extraction was performed by a single author (MH).

Data analysis and interpretation

Study characteristics including design, sample size, participant demographics, and intervention characteristics were extracted. The data are presented as descriptive statistics. Primary and secondary broad domains measured, and associated instruments used in the included studies, relevant to SSc-DUs were recorded. All the authors participated in identifying the overarching disease-related domains.

Results

Study selection

The study selection process is depicted in [Fig. 1](#). 4869 records were identified before duplicates (n=1126) were removed. Of the remaining 3743 records, 123 were eligible for full-text screening and 40 were included in the final analysis. The three most common reasons for exclusion of full texts were wrong study design as specified in the eligibility criteria (n=36), abstract only (n=17), and insufficient sample size (n=14).

Study characteristics

Study characteristics are presented in [Table 1](#), including study design, intervention, comparator (where applicable), and sample sizes. The studies were published between 1985 and 2020, with the majority (n=30) published after 2010. Sample sizes varied widely in both the intervention (8 to 1439) and comparator (6 to 186) groups. Most studies were randomized controlled trials (n=13), or prospective (n=12)/retrospective (n=8) observational studies. Active interventions included oral/intravenous drug therapies (n=25), topical/local treatments (n=4), and surgical intervention (n=2).

The broad domains used to assess SSc-DUs are presented in [Table 2](#). These broad domains can be grouped under three main themes: 'DU burden', 'DU impact', and 'Special tests' [28,29]. The broad domains and instruments used to assess these are presented as **Supplementary Table 1**.

Digital ulcer burden

DU burden encompasses the number (count), healing (i.e., ulcers which take longer to heal are more burdensome), and global burden and impact of DUs. DU burden was assessed in approximately half the studies, as assessed by DU count/number (n=23) or DU improvement (n=20). DU complications such as infection or need for surgical debridement were assessed in approximately one-sixth of studies (n=7). Two studies utilised a global assessment of DUs.




Fig. 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow diagram detailing the search and study selection process.

Impact of digital ulcers

Function was assessed in one-quarter ($n=10$) of studies. Pain directly attributed to ulcers was examined in several studies ($n=6$). Health-related quality of life was assessed in four studies.

Special tests

Specialized testing including objective assessment of the microvasculature of tissues adjacent to digital ulcers using laser-based techniques was included in four studies. One study examined DU histopathology, utilizing ulcer biomarkers (i.e., ulcer wound biopsy and assessment of vascular biomarkers).

Discussion

This scoping review identified three broad-ranging themes comprising the broad domains considered important by investigators when studying SSc-DUs: 'DU Burden', 'DU function', and 'Special tests'. However, there is currently significant heterogeneity and a lack of consensus regarding the broad domains that have been used to study SSc-DUs.

DU burden was the domain most commonly reported, and predominantly related to assessment of DU count/number ($n=23$) or DU improvement ($n=20$). Although DU complications can significantly impact on healing and may require surgical intervention [9,30], these

important aspects were only reported in seven studies. The overall (global) impact of DUs in patients with SSc has been little studied, with only two studies incorporating clinician global assessment, and only one which incorporated patient global assessment (**Supplementary Table 1**).

The functional impact of DUs was assessed in one-quarter of studies ($n=10$). SSc-DUs, especially when severe, have a major impact on the lives of those affected, including impairment of physical and social activity, emotional impact, and effects on personal relationships [19]. Although pain is a cardinal feature of SSc-DUs and patients use a broad range of narrative devices to describe this symptom [19,20], only six studies assessed ulcer-related pain. Furthermore, the lived patient experience of DU pain is complex [18–20]. For example, our previous qualitative research identified five narrative devices: 'Words to express DU-associated pain', 'Descriptions of physical and psychological reactions to pain', 'Comparisons with other painful events', 'Descriptions of factors that exacerbate pain', and 'Descriptions of strategies for coping with the pain' [20]. Patients with SSc-DUs also live with significant anxiety and uncertainty and make many adaptations to try and prevent and/or mitigate future DU episodes [19]; however, this has not been specifically captured in ulcer-related studies to date.

Investigators have utilized specialized tests, including non-invasive imaging to study the pathogenesis of SSc-DU and assess response to intervention; however, there is currently significant heterogeneity in approaches to these issues and no standardized approach for utilizing such tools has emerged to date [31]. Although the pathobiology of DUs

Table 1
Characteristics of studies of digital ulcers in systemic sclerosis included in the scoping review.

First author	Year	Study design	Intervention	Comparator	Sample size *	
					Intervention	Comparator
Mohrland [32]	1985	Randomized controlled trial	Prostaglandin E1	Placebo	16	15
Williams [33]	1985	Randomized controlled trial	Topical dimethyl sulfoxide	Topical normal saline	53	31
Wigley [34]	1992	Randomized controlled trial	Iloprost	Placebo	18	17
Vayssairat [35]	1999	Randomized controlled trial	Beraprost sodium	Placebo	55	52
Bettoni [36]	2002	Prospective cohort	Iloprost		30	
Korn [37]	2004	Randomized controlled trial	Bosentan	Placebo	79	43
Gore [38]	2005	Retrospective cohort	Sildenafil		10	
Abou-Raya [39]	2008	Randomized controlled trial	Atorvastatin	Placebo + healthy volunteers	56	28
Rosato [40]	2009	Prospective cohort	N-acetylcysteine		50	
Tsifetaki [41]	2009	Prospective cohort	Bosentan		30	
Giuggioli [42]	2010	Uncontrolled trial	Oxycodone		29	
Mouthon [43]	2010	Prospective cohort	None		213	
Zelenietz [44]	2010	Retrospective analysis of Randomized control trial	Bosentan	Placebo	176	133
Bérezné [45]	2011	Prospective cohort			189	
Matucci-Cerinic [46]	2011	Randomized controlled trial	Bosentan	Placebo	98	90
Roman Ivorra [47]	2011	Retrospective cohort	Bosentan		67	
Cozzi [48]	2013	Retrospective cohort	Bosentan	Matched control group	30	30
Ennis [49]	2013	Prospective cohort			148	
Agard [50]	2014	Retrospective cohort	Bosentan		89	
Chung [51]	2014	Prospective, open-label	Ambrisentan		20	
Mouthon [52]	2014	Prospective cohort	None		190	
Barsotti [53]	2015	Retrospective cohort	Allogenic skin grafting		43	
Meijs [54]	2015	Prospective cohort	Bosentan	Healthy controls	52	51
Ruaro [55]	2015	Imaging study	None		20	
Shah [56]	2016	Retrospective cohort	Treprostinil		51	
De Cata [57]	2016	Retrospective cohort	Iloprost and bosentan		34	
Hachulla [58]	2016	Randomized controlled trial	Sildenafil	Placebo	42	42
Khanna [59]	2016	Randomized controlled trial	Macitentan	Placebo	368	186
Küçükşahin [60]	2016	Prospective study	Bosentan		30	
Matucci-Cerinic [17]	2016	Prospective cohort	None		1459	
Hamaguchi [61]	2017	Prospective cohort	Bosentan		28	
Hughes [62]	2017	Crossover study	Topical glyceryl trinitrate	Topical placebo ointment	16	
Motegi [63]	2017	Prospective single-blind controlled trial **	Botulinum toxin B injection	No treatment	37	8
Seibold [64]	2017	Randomized controlled trial	Treprostinil	Placebo	72	76
Hassanien [65]	2018	Randomized controlled trial	Topical oxygen-ozone + calcium channel blocker	Calcium channel blocker	25	25
Simpson [66]	2018	Cross-sectional study			36	
Del Papa [67]	2019	Randomized controlled trial	Regional grafting of autologous adipose tissue	Sham procedure	25	13
Gualdi [68]	2019	Retrospective cohort	Hyaluronic acid-based wound dressing		79	
Nagaraja [69]	2019	Randomized controlled trial	Riociguat	Placebo	9	8
Barsotti [70]	2020	Cross-sectional study			31	

* Number of patients with SSc included in the study.

** Single-blinded study – patients.

Table 2
Domains used in clinical research studying digital ulcers in systemic sclerosis.

Broad domain	Target domain (number of studies)
Burden of Digital Ulcers	Digital ulcer count/number [23] Digital ulcer improvement [20] Digital ulcer complications [7] Global digital ulcer assessment [2]
Impact of Digital Ulcers	Function [10] Pain [6] Health-related quality of life [4]
Special Tests	Microvascular assessment [4] Histopathology [1]

in SSc is complex, ischemia is believed to drive pathogenesis and significantly delay or impair ulcer healing. Objective microvascular assessment was performed in four studies. Laser-derived imaging methods are considered promising potential surrogates for clinical trials of SSc-DUs, and could support early-phase proof of concept studies before larger confirmatory trials are undertaken [31]. One study utilized

DU histopathology; however, this required repeated invasive ulcer biopsies. The potential role of circulating vascular biomarkers should also be examined.

Although this review benefited from a comprehensive study design, there are some limitations to consider. This review incorporated broad inclusion criteria and used multiple databases. However, because only articles written in English, and those with a minimum number of participants were considered for inclusion, it is possible that other relevant domains could have been missed. A specific definition for DUs was not required to be adhered to, which could be important considering the poor reliability for identifying DUS reported between rheumatologists with an interest in SSc and patients themselves [5,22,24]. There was no restriction to specific ulcer locations, and some studies were confined to study of fingertip and ‘non-extensor aspect’ ulcers. Most studies related to oral or intravenous drug therapies, with only a limited number of studies on topical/local treatments and surgical interventions, reflecting the lack of reports of these treatment modalities.

This scoping review identified the spectrum of relevant outcome broad domains in the study of SSc-DU. Next steps in the project include

achieving further consensus among stakeholders and voting on a core disease domain set per the OMERACT framework.

In conclusion, this scoping review identified a broad range of disease-related domains for studying SSC-DUs, including in RCTs. These results will inform the OMERACT Vascular Disease in Systemic Sclerosis Working Group in the development of a core set of disease domains to assess the impact of SSC-DUs.

Declaration of Competing Interest

Dr Hughes reports speaking fees from Actelion Pharmaceuticals, Eli Lilly, Janssen, and Pfizer. Research funding from Janssen. Member of a Data and Safety Monitoring Board for Certa Therapeutics.

Dr. Pauling reports grants and personal fees from Janssen, outside the submitted work; Dr Pauling reports personal fees from Astra Zeneca, Boehringer Ingelheim, Sojournix Pharma and Permeatus, Inc.

Dr Proudman reports receiving funds for the following activities: advisory board: Boehringer-Ingelheim, Janssen, Gossamer. Research Support: Janssen.

Dr. Merkel reports receiving funds for the following activities in the past 2 years: Consulting: AbbVie, AstraZeneca, Boehringer-Ingelheim, Bristol-Myers Squibb, Cabaletta, ChemoCentryx, CSL Behring, Dynacure, EMDSerono, Forbius, Genentech/Roche, GlaxoSmithKline, InflaRx, Janssen, Jubilant, Kiniksa, Kyverna, Magenta, MiroBio, Mitsubishi, Neutrolis, Novartis, NS Pharma, Pfizer, Q32, Regeneron, Sparrow, Takeda, Vistera. Research Support: AbbVie, AstraZeneca, Boehringer-Ingelheim, Bristol-Myers Squibb, ChemoCentryx, Eicos, Electra, Forbius, Genentech/Roche, GlaxoSmithKline, InflaRx, Sanofi, Star, Takeda. Stock options: Kyverna. Royalties: UpToDate.

Professor Herrick reports research funding from Gesynta Pharma, consultancies with Arena, Boehringer-Ingelheim, Camurus, CSL Behring, Galderma, Gesynta Pharma and speaker's fees from Janssen.

Acknowledgement

This work was supported in part by the NIHR Manchester Biomedical Research Centre.

Funding sources

None.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.semarthrit.2023.152220](https://doi.org/10.1016/j.semarthrit.2023.152220).

References

- Steen V, Denton CP, Pope JE, Matucci-Cerinic M. Digital ulcers: overt vascular disease in systemic sclerosis. *Rheumatology (Oxford)* 2009;48(Suppl 3):iii19–24.
- Hughes M, Herrick AL. Digital ulcers in systemic sclerosis. *Rheumatology (Oxford)* 2017;56(1):14–25.
- Hughes M, Allanore Y, Chung L, Pauling JD, Denton CP, Matucci-Cerinic M, et al. Raynaud's Phenomenon and Digital Ulcers in Systemic Sclerosis. *Nat Rev Rheumatol* 2020;4(4):208–21.
- Suliman YA, Bruni C, Johnson SR, Praino E, Alemam M, Borazan N, et al. Defining Skin Ulcers in Systemic Sclerosis: Systematic Literature Review and Proposed World Scleroderma Foundation (WSF) Definition. *J Scleroderma Relat Disord* 2017;2(2):115–20.
- Hughes M, Tracey A, Bhushan M, Chakravarty K, Denton CP, Dubey S, et al. Reliability of digital ulcer definitions as proposed by the UK Scleroderma Study Group: A challenge for clinical trial design. *J Scleroderma Relat Disord* 2018;3(2):170–4.
- Hachulla E, Cleron P, Launay D, Lambert M, Morell-Dubois S, Queyrel V, et al. Natural history of ischemic digital ulcers in systemic sclerosis: Single-center retrospective longitudinal study. *J Rheumatol* 2007;34(12):2423–30.
- Khimdas S, Harding S, Bonner A, Zummer B, Baron M, Pope J. Associations with digital ulcers in a large cohort of systemic sclerosis: Results from the canadian scleroderma research group registry. *Arthritis Care Res* 2011;63(1):142–9.
- Hughes M, Pauling JD, Jones J, Denton CP, Domsic RT, Frech TM, et al. Patient experiences of digital ulcer development and evolution in systemic sclerosis. *Rheumatol* 2020;59(8):2156–8.
- Amanzi L, Braschi F, Fiori G, Galluccio F, Miniati I, Conforti D, et al. Digital ulcers in scleroderma: staging, characteristics and sub-setting through observation of 1614 digital lesions. *Rheumatology (Oxford)* 2010;49(7):1374–82.
- Bartoli F, Fiori G, Braschi F, Amanzi L, Bruni C, Blagojevic J, et al. Calcinosis in systemic sclerosis: subsets, distribution and complications. *Rheumatology (Oxford)* 2016;55(9):1610–4.
- Hughes M, Herrick AL. Diagnosis and management of systemic sclerosis-related calcinosis. *Expert Rev Clin Immunol* 2023;19(1):45–54.
- Giuggioli D, Manfredi A, Colaci M, Lumetti F, Ferri C. Osteomyelitis complicating scleroderma digital ulcers. *Clin Rheumatol* 2013;32(5):623–7.
- Allanore Y, Denton CP, Krieg T, Cornelisse P, Rosenberg D, Schwierin B, et al. Clinical characteristics and predictors of gangrene in patients with systemic sclerosis and digital ulcers in the Digital Ulcer Outcome Registry: a prospective, observational cohort. *Ann Rheum Dis* 2016;75(9):1736–40.
- Bobea C, Tatu AL, Craescu M, Solovăstru L. Dynamics of digital ulcers in systemic sclerosis. *Exp Ther Med* 2020;20(1):61–7.
- Bruni C, Guiducci S, Bellando-Randone S, Lepri G, Braschi F, Fiori G, et al. Digital ulcers as a sentinel sign for early internal organ involvement in very early systemic sclerosis. *Rheumatology (Oxford)* 2015;54(1):72–6. Jan.
- Mihai C, Landewé R, van der Heijde D, Walker UA, Constantin PI, Gherghe AM, et al. Digital ulcers predict a worse disease course in patients with systemic sclerosis. *Ann Rheum Dis* 2016;75(4):681–6.
- Matucci-Cerinic M, Krieg T, Guillevin L, Schwierin B, Rosenberg D, Cornelisse P, et al. Elucidating the burden of recurrent and chronic digital ulcers in systemic sclerosis: long-term results from the DUO Registry. *Ann Rheum Dis* 2016;75(10):1770–6.
- Hughes M, Pauling JD. Exploring the patient experience of digital ulcers in systemic sclerosis. *Semin Arthritis Rheum* 2019;48(5):888–94.
- Hughes M, Pauling JD, Jones J, Denton CP, Domsic RT, Frech TM, et al. Multicenter Qualitative Study Exploring the Patient Experience of Digital Ulcers in Systemic Sclerosis. *Arthritis Care Res (Hoboken)* 2020;72(5):723–33.
- Jones J, Hughes M, Pauling J, Goberman-Hill R, Moore AJ. What narrative devices do people with systemic sclerosis use to describe the experience of pain from digital ulcers: a multicentre focus group study at UK scleroderma centres. *BMJ Open* 2020;10(6):e037568.
- Morrisroe K, Stevens W, Sahhar J, Ngian GS, Ferdowsi N, Hill CL, et al. Digital ulcers in systemic sclerosis: Their epidemiology, clinical characteristics, and associated clinical and economic burden. *Arthritis Res Ther* 2019;21(1):1–12.
- Herrick AL, Roberts C, Tracey A, Silman A, Anderson M, Goodfield M, et al. Lack of agreement between rheumatologists in defining digital ulceration in systemic sclerosis. *Arthritis Rheum* 2009;60(3):878–82. Mar.
- Baron M, Chung L, Gyger G, Hummers L, Khanna D. Consensus opinion of a North American Working Group regarding the classification of digital ulcers in systemic sclerosis. *Clin Rheumatol* 2014;33(2):207–14.
- Hughes M, Roberts C, Tracey A, Dinsdale G, Murray A, Herrick ALL. Does the Clinical Context Improve the Reliability of Rheumatologists Grading Digital Ulcers in Systemic Sclerosis? *Arthritis Care Res* 2016;68(9):1340–5.
- Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol* 2005;8(1):19–32.
- Maltez N, Hughes M, Brown E, Hickey V, Park H, Shea B, et al. Developing a core set of outcome measure domains to study Raynaud's phenomenon and digital ulcers in systemic sclerosis: Report from OMERACT 2020. *Semin Arthritis Rheum* 2021;51(3):640–3.
- Beaton D, Maxwell L, Grosskleg S, Shea B, Tugwell B, editors. *The OMERACT Handbook Version 2.1*. 2021.
- Beaton LJ, Beaton DE, Shea BJ, Wells GA, Boers M, Grosskleg S, et al. Core Domain Set Selection According to OMERACT Filter 2.1: The OMERACT Methodology. *J Rheumatol* 2019;46(8):1014–20.
- Boers M, Beaton DE, Shea BJ, Maxwell LJ, Bartlett SJ, Bingham 3rd CO, et al. OMERACT Filter 2.1: Elaboration of the Conceptual Framework for Outcome Measurement in Health Intervention Studies. *J Rheumatol* 2019;46(8):1021–7.
- Hughes M, Allanore Y, El Aoufy K, Denton CP, Khanna D, Krieg T, et al. A Practical Approach to the Management of Digital Ulcers in Patients With Systemic Sclerosis: A Narrative Review. *JAMA Dermatology* 2021;157(7):851–8.
- Pauling JD, Hackett N, Guida A, Merkel PA. Performance of laser-derived imaging for assessing digital perfusion in clinical trials of systemic sclerosis-related digital vasculopathy: A systematic literature review. *Semin Arthritis Rheum* 2020;50(5):1114–30.
- Mohrland JS, Porter JM, Smith EA, Belch J, Simms MH. A multiclinic, placebo-controlled, double-blind study of prostaglandin E1 in Raynaud's syndrome. *Ann Rheum Dis* 1985;44(11):754–60.
- Williams HJ, Furst DE, Dahl SL, Steen VD, Marks C, Alpert EJ, et al. Double-blind, multicenter controlled trial comparing topical dimethyl sulfoxide and normal saline for treatment of hand ulcers in patients with systemic sclerosis. *Arthritis Rheum* 1985;28(3):308–14.
- Wigley FM, Seibold JR, Wise RA, McCloskey DA, Dole WP. Intravenous iloprost treatment of Raynaud's phenomenon and ischemic ulcers secondary to systemic sclerosis. *J Rheumatol* 1992;19(9):1407–14.
- Vayssairat M. Preventive effect of an oral prostacyclin analog, beraprost sodium, on digital necrosis in systemic sclerosis. French Microcirculation Society Multicenter Group for the Study of Vascular Acrosyndromes. *J Rheumatol* 1999;26(10):2173–8.

- [36] Bettoni L, Geri A, Airò P, Danieli E, Cavazzana I, Antonioli C, et al. Systemic sclerosis therapy with iloprost: a prospective observational study of 30 patients treated for a median of 3 years. *Clin Rheumatol* 2002;21(3):244–50.
- [37] Korn JH, Mayes M, Matucci Cerinic M, Rainisio M, Pope J, Hachulla E, et al. Digital ulcers in systemic sclerosis: prevention by treatment with bosentan, an oral endothelin receptor antagonist. *Arthritis Rheum* 2004;50(12):3985–93.
- [38] Gore J, Silver R. Oral sildenafil for the treatment of Raynaud's phenomenon and digital ulcers secondary to systemic sclerosis. *Ann Rheum Dis* 2005;64(9):1387.
- [39] Abou-Raya A, Abou-Raya S, Helmi M. Statins: potentially useful in therapy of systemic sclerosis-related Raynaud's phenomenon and digital ulcers. *J Rheumatol* 2008;35(9):1801–8.
- [40] Rosato E, Borghese F, Pisarri S, Salsano F. The treatment with N-acetylcysteine of Raynaud's phenomenon and ischemic ulcers therapy in scleroderma patients: a prospective observational study of 50 patients. *Clin Rheumatol* 2009;28(12):1379–84.
- [41] Tsifetaki N, Botzorris V, Alamanos Y, Argyriou E, Zioga A, Drosos AA. Bosentan for digital ulcers in patients with systemic sclerosis: a prospective 3-year followup study. *J Rheumatol* 2009;36(7):1550–2.
- [42] Giuggioli D, Manfredi A, Colaci M, Ferri C. Oxycodone in the long-term treatment of chronic pain related to scleroderma skin ulcers. *Pain Med* 2010;11(10):1500–3.
- [43] Mouthon L, Mestre-Stanislas C, Bérezné A, Rannou F, Guilpain P, Revel M, et al. Impact of digital ulcers on disability and health-related quality of life in systemic sclerosis. *Ann Rheum Dis* 2010;69(1):214–7.
- [44] Zelenietz C, Pope J. Differences in disability as measured by the Health Assessment Questionnaire between patients with and without digital ulcers in systemic sclerosis: a post hoc analysis of pooled data from two randomised controlled trials in digital ulcers using bosentan. *Ann Rheum Dis*;69(11):2055–6.
- [45] Bérezné A, Seror R, Morell-Dubois S, de Menthon M, Fois E, Dzeing-Ella A, et al. Impact of systemic sclerosis on occupational and professional activity with attention to patients with digital ulcers. *Arthritis Care Res* 2011;63(2):277–85.
- [46] Matucci-Cerinic M, Denton CP, Furst DE, Mayes MD, Hsu VM, Carpentier P, et al. Bosentan treatment of digital ulcers related to systemic sclerosis: results from the RAPIDS-2 randomised, double-blind, placebo-controlled trial. *Ann Rheum Dis* 2011;70(1):32–8.
- [47] Roman Ivorra JA, Simeon CP, Alegre Sancho JJ, Egurbide MV, Castillo MJ, Lloria X, et al. Bosentan in clinical practice for treating digital and other ischemic ulcers in Spanish patients with systemic sclerosis: IBER-DU cohort study. *J Rheumatol* 2011;38(8):1631–5.
- [48] Cozzi F, Pigatto E, Rizzo M, Favaro M, Zanatta E, Cardarelli S, et al. Low occurrence of digital ulcers in scleroderma patients treated with bosentan for pulmonary arterial hypertension: a retrospective case-control study. *Clin Rheumatol* 2013;32(5):679–83.
- [49] Ennis H, Vail A, Wragg E, Taylor Moore T, Murray A, et al. A prospective study of systemic sclerosis-related digital ulcers: prevalence, location, and functional impact. *Scand J Rheumatol* 2013;42(6):483–6.
- [50] Agard C, Carpentier PH, Mouthon L, Clerson P, Gressin V, Bérezné A, et al. Use of bosentan for digital ulcers related to systemic sclerosis: a real-life retrospective French study of 89 patients treated since specific approval. *Scand J Rheumatol* 2014;43(5):398–402.
- [51] Chung L, Ball K, Yaqub A, Lingala B, Fiorentino D. Effect of the endothelin type A-selective endothelin receptor antagonist ambrisentan on digital ulcers in patients with systemic sclerosis: results of a prospective pilot study. *J Am Acad Dermatol* 2014;71(2):400–1.
- [52] Mouthon L, Carpentier PH, Lok C, Clerson P, Gressin V, Hachulla E, et al. Ischemic digital ulcers affect hand disability and pain in systemic sclerosis. *J Rheumatol* 2014;41(7):1317–23.
- [53] Barsotti S, Mattaliano V, d'Ascanio A, Mosti G, Della Rossa A, Mattaliano C, et al. Systemic sclerosis chronic ulcers: preliminary results of treatment with allogenic skin grafting in a cohort of Italian patients. *Int Wound J* 2016;13(5):1050–1.
- [54] Meijs J, Voskuyl AE, Bloemsaat-Minekus JPJ, Vonk MC. Blood flow in the hands of a predefined homogeneous systemic sclerosis population: the presence of digital ulcers and the improvement with bosentan. *Rheumatology (Oxford)* 2015;54(2):262–9.
- [55] Ruaro B, Sulli A, Smith V, Paolino S, Pizzorni C, Cutolo M. Short-term follow-up of digital ulcers by laser speckle contrast analysis in systemic sclerosis patients. *Microvasc Res* 2015;101:82–5.
- [56] Shah AA, Schiopu E, Chatterjee S, Csuka ME, Frech T, Goldberg A, et al. The Recurrence of Digital Ulcers in Patients with Systemic Sclerosis after Discontinuation of Oral Treprostinil. *J Rheumatol* 2016;43(9):1665–71.
- [57] De Cata A, Inglese M, Molinaro F, De Cosmo S, Rubino R, Bernal M, et al. Digital ulcers in scleroderma patients: A retrospective observational study. *Int J Immunopathol Pharmacol* 2016;29(2):180–7.
- [58] Hachulla E, Hatron PY, Carpentier P, Agard C, Chatelus E, Jegou P, et al. Efficacy of sildenafil on ischaemic digital ulcer healing in systemic sclerosis: the placebo-controlled SEDUCE study. *Ann Rheum Dis* 2016;75(6):1009–15.
- [59] Khanna D, Denton CP, Merkel PA, Krieg TM, Le Brun FO, Marr A, et al. Effect of macitentan on the development of new ischemic digital ulcers in patients with systemic sclerosis: Dual-1 and Dual-2 randomized clinical trials. *JAMA* 2016;315(18):1975–88.
- [60] Küçükşahin O, Yildizgören MT, Gerede DM, Maraş Y, Erten Ş. Bosentan For Digital Ulcers in Patients With Systemic Sclerosis: Single Center Experience. *Arch Rheumatol* 2016;31(3):229–33.
- [61] Hamaguchi Y, Sumida T, Kawaguchi Y, Ihn H, Tanaka S, Asano Y, et al. Safety and tolerability of bosentan for digital ulcers in Japanese patients with systemic sclerosis: Prospective, multicenter, open-label study. *J Dermatol* 2017;44(1):13–7.
- [62] Hughes M, Moore T, Manning J, Wilkinson J, Dinsdale G, Roberts C, et al. Reduced perfusion in systemic sclerosis digital ulcers (both fingertip and extensor) can be increased by topical application of glyceryl trinitrate. *Microvasc Res* 2017;111:32–6.
- [63] Motegi SI, Uehara A, Yamada K, Sekiguchi A, Fujiwara C, Toki S, et al. Efficacy of botulinum toxin B injection for Raynaud's phenomenon and digital ulcers in patients with systemic sclerosis. *Acta Derm Venereol* 2017;97(7):843–50.
- [64] Seibold JR, Wigley FM, Schiopu E, Denton CP, Silver RM, Steen VD, et al. Digital ulcers in SSc treated with oral treprostinil: a randomized, double-blind, placebo-controlled study with open-label follow-up. *J Scleroderma Relat Disord* 2017;2(1):42–9.
- [65] Hassanien M, Rashad S, Mohamed N, Elawamy A, Ghaly MS. Non-invasive Oxygen-Ozone therapy in treating digital ulcers of patients with systemic sclerosis. *Acta Reumatol Port* 2018;2018(3):210–6.
- [66] Simpson V, Hughes M, Wilkinson J, Herrick AL, Dinsdale G. Quantifying Digital Ulcers in Systemic Sclerosis: Reliability of Computer-Assisted Planimetry in Measuring Lesion Size. *Arthritis Care Res* 2018;70(3):486–90.
- [67] Del Papa N, Di Luca G, Andracco R, Zaccara E, Maglione W, Pignataro F, et al. Regional grafting of autologous adipose tissue is effective in inducing prompt healing of indolent digital ulcers in patients with systemic sclerosis: Results of a monocentric randomized controlled study. *Arthritis Res Ther* 2019;21(1):1–11.
- [68] Gualdi G, Monari P, Cammalleri D, Pelizzari L, Pinton PC. Hyaluronic Acid-based Products are Strictly Contraindicated in Scleroderma-related Skin Ulcers. *Wounds* 2019;31(3):81–4.
- [69] Nagaraja V, Spino C, Bush E, Tsou PS, Domsic RT, Lafyatis R, et al. A multicenter randomized, double-blind, placebo-controlled pilot study to assess the efficacy and safety of riociguat in systemic sclerosis-associated digital ulcers. *Arthritis Res Ther* 2019;21(1):1–14.
- [70] Barsotti S, d'Ascanio A, Valentina V, Chiara S, Silvia B, Laura A, et al. Is there a role for laser speckle contrast analysis (LASCA) in predicting the outcome of digital ulcers in patients with systemic sclerosis? *Clin Rheumatol* 2020;39(1):69–75.